

The failure of Selman's Theorem for hyperenumeration reducibility

Josiah Jacobsen-Grocott

University of Wisconsin—Madison
Partially supported by NSF Grant No. DMS-2053848

Specialty Exam, February, 2022

Table of contents

- 1 Enumeration reducibility and Selman's Theorem
- 2 Hyperenumeration reducibility
- 3 E-pointed trees
- 4 Computability in $L_{\omega_1^{CK}}$
- 5 A non-hypertotal e-pointed tree

Table of contents

- 1 Enumeration reducibility and Selman's Theorem
- 2 Hyperenumeration reducibility
- 3 E-pointed trees
- 4 Computability in $L_{\omega_1}^{CK}$
- 5 A non-hypertotal e-pointed tree

Definition

For two sets $A, B \subseteq \omega$ we say that $A \leq_e B$ if there is a c.e. set W such that:

$$x \in A \iff \exists \langle x, u \rangle \in W [D_u \subseteq B]$$

where $(D_u)_{u \in \omega}$ is listing of all finite sets by strong indices.

- From an effective listing of c.e. sets $(W_e)_{e \in \omega}$ we obtain an effective listing of enumeration operators $(\Psi_e)_{e \in \omega}$. Defined by $A = \Psi_e(B)$ if $A \leq_e B$ via W_e .
- \leq_e is a preorder and, like with Turing reducibility and the Turing degrees, we get the enumeration degrees \mathcal{D}_e .

Definition

We say that a set A is *total* if $\bar{A} \leq_e A$. We say that A is *cototal* if $A \leq_e \bar{A}$. A degree is *total* (*cototal*) if it contains a total (cototal) set.

- If A is total then $B \leq_e A$ if and only if B is c.e. in A .
- For any set A we have that $A \oplus \bar{A}$ is both total and cototal.
- The Turing degrees embed onto the total degrees via the map induced by $A \mapsto A \oplus \bar{A}$.
- The cototal degrees are a proper subclass of the enumeration degrees and the total degrees are a proper subclass of the cototal degrees.

Selman's Theorem

As we have seen, we can define Turing reducibility in terms of enumeration reducibility. Selman's theorem gives us a way of defining enumeration reducibility in terms of Turing reducibility.

Theorem (Selman's Theorem)

$A \leq_e B$ if and only if for all X if $B \leq_e X \oplus \bar{X}$ then $A \leq_e X \oplus \bar{X}$.

There is another way to define enumeration reducibility in terms of enumerations. We have that $A \leq_e B$ if every enumeration of B uniformly computes an enumeration of A . Here an enumeration of A is a total, onto function $f : \omega \rightarrow A$. In this context, Selman's theorem shows that we can drop the uniformity in the definition..

Proof of Selman's Theorem

Proof.

Suppose that $B \not\leq_e A$. We will use forcing to build an enumeration f of A that is not above B . At stage s given initial segment $\sigma_s \in \omega^{<\omega}$ we ask if there is $\tau \succeq \sigma_s$ and $n \notin B$ such that $n \in \Psi_s(\tau)$ and $\text{range}(\tau) \subseteq A$. If there is such a τ then we set $\sigma_{s+1} = \tau$. If there is no such τ then let $k = \min(A \setminus \text{range}(\sigma_s))$ and set $\sigma_{s+1} = \sigma_s \hat{\ } k$.

By construction we have that $f = \bigcup_s \sigma_s$ is an enumeration of A . Now suppose towards a contradiction that $B = \Psi_e(f)$ for some e . Then at stage e we must not have found any τ . So for all $\tau \succ \sigma_e$ with $\text{range}(\tau) \subseteq A$ we have that $\Psi_e(\tau) \subseteq B$. So as $B = \Psi_e(f)$ we have:

$$n \in B \iff \exists \tau \succ \sigma_e [\text{range}(\tau) \subseteq A]$$

Hence $B \leq_e A$. □

Table of contents

- 1 Enumeration reducibility and Selman's Theorem
- 2 Hyperenumeration reducibility
- 3 E-pointed trees
- 4 Computability in $L_{\omega_1}^{CK}$
- 5 A non-hypertotal e-pointed tree

Hyperenumeration reducibility

- Now we define hyperenumeration reducibility as introduced by Sanchis in 1978.

Definition

We say that $A \leq_{he} B$ if there is a c.e. set W such that

$$n \in A \iff \forall f \in \omega^\omega \exists u \in \omega, x \prec f[\langle n, x, u \rangle \in W \wedge D_u \subseteq B]$$

- Like with enumeration reducibility this is a preorder and the equivalence classes give us the hyperenumeration degrees \mathcal{D}_{he} .
- From an effective listing of c.e. sets $(W_e)_{e \in \omega}$ we obtain an effective listing of hyperenumeration operators $(\Gamma_e)_{e \in \omega}$.

Hypertotal degrees.

Definition

We say that a set A is *hypertotal* if $\bar{A} \leq_{he} A$. We say that A is *hypercototal* if $A \leq_{he} \bar{A}$. A degree (in either \mathcal{D}_e or \mathcal{D}_{he}) is *hypertotal* (*hypercototal*) if it contains a hypertotal (hypercototal) set.

We have a similar relationship between the hypertotal degrees and the hyperarithmetical degrees as the relationship between the total and Turing degrees.

From the definition of \leq_{he} we have that if $A \leq_{he} B$ then A is Π_1^1 in B . It is not hard to show that if A is Π_1^1 in B then $A \leq_{he} B \oplus \bar{B}$. So $A \leq_h B \iff A \oplus \bar{A} \leq_{he} B \oplus \bar{B}$. The hyperarithmetical degrees embed onto the total degrees via the map induced by $A \mapsto A \oplus \bar{A}$.

Theorem (Sanchis)

There is a hyperenumeration degree that is not hypertotal.

Sanchis proved an interesting result about the relationship between enumeration reducibility and hyperenumeration reducibility.

Theorem (Sanchis)

If $A \leq_e B$ then $A \leq_{he} B$ and $\bar{A} \leq_{he} \bar{B}$.

This means that if f is an enumeration of A then $A \oplus \bar{A} \leq_{he} f$. So when working with hyperenumeration reducibility we want a new notion of a hyperenumeration.

Hyperenumerations

Recall the definition of $A = \Gamma_e(B)$.

$$n \in A \iff \forall f \in \omega^\omega \exists u \in \omega, x \prec f[\langle n, x, u \rangle \in W_e \wedge D_u \subseteq B]$$

Now consider the tree $S_e \subseteq \omega^{<\omega}$ defined by

$$n \hat{\ } x \notin S_e \iff \exists y \preceq x, u \leq |x| [\langle n, y, u \rangle \in W_{e,|x|} \wedge D_u \subseteq B]$$

We have that $S_e \leq_T B$ and $\overline{S_e} \leq_e B$. Define $S_{e,n} = \{x : n \hat{\ } x \in S_e\}$. We have that

$$n \in A \iff S_{e,n} \text{ is well founded}$$

So $A \leq_{he} \overline{S_e}$. We call a tree which hyperenumerates A in the way that S_e does a hyperenumeration of A .

Table of contents

- 1 Enumeration reducibility and Selman's Theorem
- 2 Hyperenumeration reducibility
- 3 E-pointed trees**
- 4 Computability in $L_{\omega_1}^{CK}$
- 5 A non-hypertotal e-pointed tree

Definition

A tree T is *e-pointed* if for every path $P \in [T]$ we have that T is c.e. in P . We say T is *uniformly e-pointed* if there is a single operator Ψ_e such that for all paths $P \in [T]$ we have $T = \Psi_e(P)$.

McCarthy studied e-pointed trees in Cantor space and was able to characterize their enumeration degrees.

Theorem (McCarthy)

If $T \subseteq 2^{<\omega}$ is uniformly e-pointed then T is cototal. Furthermore for a degree $a \in \mathcal{D}_e$ the following are equivalent:

- a is cototal.
- a contains an e-pointed tree $T \subseteq 2^{<\omega}$.
- a contains a uniformly e-pointed tree $T \subseteq 2^{<\omega}$ with no dead ends.

E-pointed trees in Baire space with dead ends

In Baire space we have the following characterization in terms of hypertotal sets.

Theorem (Goh, J-G, Miller, Soskova)

If $T \subseteq \omega^{<\omega}$ is uniformly e-pointed then T is hypertotal. Furthermore for a degree $a \in \mathcal{D}_e$ (or \mathcal{D}_{he}) the following are equivalent:

- *a is hypercototal.*
- *a contains an e-pointed tree $T \subseteq \omega^{<\omega}$.*
- *a contains a uniformly e-pointed tree $T \subseteq \omega^{<\omega}$.*

E-pointed trees in Baire space without dead ends

When we consider only e-pointed trees that do not have dead ends then things become more complex

Theorem (Goh, J-G, Miller, Soskova)

There is an arithmetic set that is not enumeration equivalent to any e-pointed tree $T \subseteq \omega^{<\omega}$ without dead ends.

Theorem (Goh, J-G, Miller, Soskova)

There is a uniformly e-pointed tree $T \subseteq \omega^{<\omega}$ without dead ends that is not of cototal enumeration degree.

Question

Is there an e-pointed tree $T \subseteq \omega^{<\omega}$ without dead ends that is not enumeration equivalent to any uniformly e-pointed tree $T \subseteq \omega^{<\omega}$ without dead ends.

Theorem (J-G)

There is a uniformly e -pointed tree with no dead ends that is not hypertotal.

This leads us to a contradiction of Selman's theorem.

Corollary

There are sets A, B such that $B \not\leq_{he} A$ and for any X , if $A \leq_{he} X \oplus \bar{X}$ then $B \leq_{he} X \oplus \bar{X}$.

Connection to Selman's theorem

Corollary

There are sets A, B such that $B \not\leq_{he} A$ and for any X , if $A \leq_{he} X \oplus \bar{X}$ then $B \leq_{he} X \oplus \bar{X}$.

Proof.

We will have $A = T$ and $B = \bar{T}$ where T is a uniformly e-pointed tree with no dead ends that is not hypertotal. Suppose that T is Π_1^1 in X . Since T has no dead ends there must be a path $P \in [T]$ such that $P \leq_h X$. So $T \leq_e P$ and by previous lemma we have $\bar{T} \leq_{he} \bar{P} \leq_h X$. So we get that $\bar{T} \leq_{he} X \oplus \bar{X}$. □

Table of contents

- 1 Enumeration reducibility and Selman's Theorem
- 2 Hyperenumeration reducibility
- 3 E-pointed trees
- 4 Computability in $L_{\omega_1^{CK}}$
- 5 A non-hypertotal e-pointed tree

Admissible sets

The usual definition of a Π_1^1 set of natural numbers is a set of the form $m \in X \iff \forall f \in \omega^\omega \exists n [R(f, n, m)]$ where R is a computable relation. However admissibility gives us another definition in terms of $L_{\omega_1^{CK}}$ that is useful.

Definition

A set M is *admissible* if it is transitive, closed under union, pairing and Cartesian product as well as satisfying the following two properties:

Δ_1 -comprehension: for every Δ_1 definable class $A \subseteq M$ and set $a \in M$ the set $A \cap a \in M$.

Σ_1 -collection: for every Σ_1 definable class relation $R \subseteq M^2$ and set $a \in M$ such that $a \subseteq \text{dom}(R)$ there is $b \in M$ such that $a = R^{-1}[b]$.

- The smallest admissible set is HF the collection of hereditarily finite sets. Looking at the Δ_1 and Σ_1 subsets of HF is one notion of computability. We have that the Δ_1 subsets of HF are computable sets and the Σ_1 subsets of HF are c.e. sets.
- We generalize this to an arbitrary admissible set M by calling a set $A \subseteq M$ M -computable if it is a Δ_1 subset of M and M -c.e. if it is a Σ_1 subset of M .
- The smallest admissible set containing ω as an element is $L_{\omega_1^{CK}}$. We have that the $L_{\omega_1^{CK}}$ -c.e. subsets of ω are precisely the Π_1^1 sets.

Table of contents

- 1 Enumeration reducibility and Selman's Theorem
- 2 Hyperenumeration reducibility
- 3 E-pointed trees
- 4 Computability in $L_{\omega_1}^{CK}$
- 5 A non-hypertotal e-pointed tree

The forcing partial order

Let $\{T_\sigma : \sigma \in \omega^{<\omega}\}$ be an effective listing of all finite trees in $\omega^{<\omega}$ where for each $\sigma \in \omega^{<\omega}$ sequence $T_{\sigma \smallfrown 0}, T_{\sigma \smallfrown 1}, \dots$ lists each finite tree that contains T_σ infinitely often.

Definition

A condition p is a pair $(T^p, L^p : T^p \times T^p \rightarrow \omega_1^{CK}) \in L_{\omega_1^{CK}}$ such that:

- 1 $T^p \subseteq \omega^{<\omega}$ is a well founded tree.
- 2 For each $\sigma \in T^p$ we have that $T_\sigma \subseteq T^p$.
- 3 $L^p(\sigma, \tau) = 0$ if and only if $\sigma \in T_\tau$.
- 4 If $\rho \prec \tau$ then $L^p(\sigma, \tau) = 0$ or $L^p(\sigma, \tau) < L^p(\sigma, \rho)$.
- 5 For each $\tau \in T^p$ and $n < \omega$ the set $\{\sigma : L^p(\sigma, \tau) \leq n\}$ is finite.

For two conditions p and q we say $p \leq q$ if $T^q \preceq T^p$ and $L^q \subseteq L^p$.

Lemma

The set of conditions is $L_{\omega_1^{\text{CK}}}$ -c.e. and the relation \leq on conditions is $L_{\omega_1^{\text{CK}}}$ -computable.

Lemma

Let $A \subseteq \omega^{<\omega}$ be a set such that for all $\sigma \hat{\ } i \in A$ we have $\sigma \in T^P$ and $\{\tau : L^P(\tau, \sigma) \leq 1\} \subseteq T_{\sigma \hat{\ } i} \subseteq T^P \cup A$. For such an A we can define a condition $q = p[A]$ with $T^q = T^P \cup A$ such that q is a valid condition. If we also have that $T^P \preceq T^P \cup A$ then $q \leq p$.

Corollary

If \mathcal{G} is a sufficiently generic filter then $T^{\mathcal{G}}$ is a uniformly e -pointed tree with no dead ends.

The forcing relation

Definition

For a condition p we define $S_e^p \subseteq \omega^\omega$ to be the tree where

$$n \hat{\ } x \notin S_e^p \iff \exists y \prec x, u \leq |x| [\langle n, y, u \rangle \in W_{e,|x|} \wedge D_u \subseteq T^p]$$

For a filter \mathcal{G} we define $S_e^{\mathcal{G}} \cap_{p \in \mathcal{G}} S_e^p$.

We define $p \Vdash \text{rank}(S_{e,x}^{\mathcal{G}}) \leq \alpha$ if $\text{rank}(S_{e,x}^p) \leq \alpha$.

So by definition of Γ_e we have $\Gamma_e(T^{\mathcal{G}}) = \{n : S_{e,n}^{\mathcal{G}} \text{ is well founded}\}$.

From this definition it is clear that if $p \Vdash \text{rank}(S_{e,x}^{\mathcal{G}}) \leq \alpha$ then for any $\mathcal{G} \ni p$ we have that $\text{rank}(S_{e,x}^{\mathcal{G}}) \leq \alpha$.

Lemma

Fix a condition p . Suppose that for each $i \in \omega$, $r \leq p$ there is $q \leq r$ such that $q \Vdash \text{rank}(S_{e,x \cap i}^{\mathcal{G}}) \leq \beta$ for some $\beta < \omega_1^{\text{CK}}$ then there is $\hat{p} \leq p$ and $\alpha < \omega_1^{\text{CK}}$ such that $\hat{p} \Vdash \text{rank}(S_{e,x}^{\mathcal{G}}) \leq \alpha$.

Lemma

If for all $q \leq p$ and $\alpha < \omega_1^{\text{CK}}$ we have $q \not\Vdash \text{rank}(S_{e,x}^{\mathcal{G}}) \leq \alpha$ then $p \Vdash S_{e,x}^{\mathcal{G}}$ is ill founded. Formally, for all sufficiently generic filters $\mathcal{G} \ni p$ we have that $S_{e,x}^{\mathcal{G}}$ contains an infinite path.

Main result

Theorem (J-G)

There is a uniformly e -pointed tree in $T^{\mathcal{G}} \subseteq \omega^{<\omega}$ with no dead ends such that $T^{\mathcal{G}}$ is not hypertotal.

Proof.

We say $p \Vdash \overline{T^{\mathcal{G}}} \neq \Gamma_e(T^{\mathcal{G}})$ if there is $\sigma \in T^P$ and $\alpha < \omega_1^{CK}$ such that $p \Vdash \text{rank}(S_{e, \langle \sigma \rangle}^{\mathcal{G}}) \leq \alpha$, or if there is $\sigma \notin T^P$ such that the initial segment of σ in T^P is not a leaf and $p \Vdash S_{e, \langle \sigma \rangle}^{\mathcal{G}}$ is ill founded. To show that $T^{\mathcal{G}}$ is not hypertotal it is enough for us to show that the sets $\{p : p \Vdash \overline{T^{\mathcal{G}}} \neq \Gamma_e(T^{\mathcal{G}})\}$ are dense for each e .

Proof continued.

Suppose towards a contradiction, that $\{p : p \Vdash \overline{T^{\mathcal{G}}} \neq \Gamma_e(T^{\mathcal{G}})\}$ is not dense. Let p be such that for all $q \leq p$ we have $q \not\Vdash \overline{T^{\mathcal{G}}} \neq \Gamma_e(T^{\mathcal{G}})$. Consider some leaf $\sigma \in T^p$ and let i, j be such that $T_{\sigma \frown i} = T_{\sigma \frown j} = \{\rho : L^p(\rho, \sigma) \leq 1\}$. Now consider $q = p[\{\sigma \frown i\}]$; this is well defined by previous lemma. By assumption on p we have that $q \not\Vdash S_{e, \langle \sigma \frown j \rangle}^{\mathcal{G}}$ is ill founded, so by previous lemma there is $r \leq q, \alpha < \omega_1^{\text{CK}}$ such that $r \Vdash \text{rank}(S_{e, \langle \sigma \frown j \rangle}^{\mathcal{G}}) \leq \alpha$. Now consider $r' = r[\{\sigma \frown j\}]$. Since $\sigma \frown i \in T^r$ we have $\{\rho : L^r(\rho, \sigma) \leq 1\} \subseteq T_{\sigma \frown i} = T_{\sigma \frown j}$ and thus the condition r' is a valid condition. Since $r \leq p$ and σ is a leaf in T^p we have that $r' \leq p$. But we have $S_e^r \supseteq S_e^{r'}$ so $r' \Vdash \text{rank}(S_{e, \langle \sigma \frown j \rangle}^{\mathcal{G}}) \leq \alpha$ a contradiction. So we have that the set $\{p : p \Vdash \overline{T^{\mathcal{G}}} \neq \Gamma_e(T^{\mathcal{G}})\}$ is dense.

So for sufficiently generic \mathcal{G} we have that $T^{\mathcal{G}}$ is uniformly e -pointed without dead ends and for all e we have $\overline{T^{\mathcal{G}}} \neq \Gamma_e(T^{\mathcal{G}})$, and thus $\overline{T^{\mathcal{G}}} \not\leq_{he} T^{\mathcal{G}}$. \square

Thank you

Thank You